

PATIENT INFORMATION

Date:	Patient Name:	DOB:
ICD-10 code (required):	ICD-10 description:	
<input type="checkbox"/> NKDA Allergies:	Weight lbs/kg:	
Patient Status: <input type="checkbox"/> New to Therapy	<input type="checkbox"/> Continuing Therapy	Next Due Date (if applicable):

PROVIDER INFORMATION

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

NURSING

- TB status and date (results) _____
 - Provide nursing care per IVX Nursing Procedures, including reaction management and post-procedure observation
- NOTE:** IVX Adverse Reaction Management Protocol available for review at www.ivxhealth.com/forms (version 09.07.2021)

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
 - cetirizine (Zyrtec) 10mg PO
 - loratadine (Claritin) 10mg PO
 - diphenhydramine (Benadryl) 25mg / 50mg PO / IV
 - methylprednisolone (Solu-Medrol) 40mg / 125mg IV
 - hydrocortisone (Solu-Cortef) 100mg IV
 - Other: _____
- Dose: _____ Route: _____
- Frequency: _____

THERAPY ADMINISTRATION

- Abatacept** (Orencia) in 100ml 0.9% sodium chloride, intravenous infusion over 30 minutes (use in line filter 0.2 to 1.2 micron)
 - Dose: 500mg / 750mg / 1000mg / _____mg
 - Frequency: induction: week 0, 2, and 4, then every 4 weeks / maintenance: every 4 weeks / other: _____
- Route: intravenous
- Infuse over 30 minutes
- Remove equal volume from bag prior to adding medication
- Flush with 0.9% sodium chloride at infusion completion
- abatacept** (Orencia) injection
 - Dose: 50mg/ 87.5mg/ 125mg
 - Frequency: weekly / other: _____
 - Route: subcutaneous
- Patient is required to stay for 30-minute observation period
- Patient is NOT required to stay for observation time
- Refills: Zero / for 12 months / _____ (if not indicated order will expire one year from date signed)

SPECIAL INSTRUCTIONS

Screen for latent TB infection prior to initiating therapy. Patients testing positive should be treated prior to initiating ORENCIA.

Provider Name (Print) _____ Provider Signature _____ Date _____

Email ivxintake@ivxhealth.com or fax this form, insurance card (both sides), demographics, recent H&P, labs, and supporting clinicals to:

<input type="checkbox"/> BAY AREA: 844-899-0275	<input type="checkbox"/> COLUMBUS: 844-627-2675	<input type="checkbox"/> JACKSONVILLE: 904-212-2338	<input type="checkbox"/> PHILADELPHIA: 844-820-9641	<input type="checkbox"/> WEST TN: 888-615-1445
<input type="checkbox"/> CINCINNATI: 844-946-0868	<input type="checkbox"/> HARRISBURG: 844-859-4235	<input type="checkbox"/> KANSAS CITY: 844-900-1292	<input type="checkbox"/> TAMPA: 844-946-0849	<input type="checkbox"/> MIDDLE TN: 888-615-1445
<input type="checkbox"/> CHICAGO: 312-253-7244	<input type="checkbox"/> INDIANAPOLIS: 844-983-2028	<input type="checkbox"/> ORLANDO: 844-946-0867	<input type="checkbox"/> WEST FLORIDA: 844-946-0849	<input type="checkbox"/> EAST TN: 888-615-1445