

Certolizumab (Cimzia)



Provider Order Form rev. 3/25/2022

PATIENT INFORMATION

Date:	Patient Name:	DOB:
ICD-10 code (required):	ICD-10 description:	
<input type="checkbox"/> NKDA Allergies:	Weight lbs/kg:	
Patient Status: <input type="checkbox"/> New to Therapy <input type="checkbox"/> Continuing Therapy	Next Due Date (if applicable):	

PROVIDER INFORMATION

Referral Coordinator Name:	Referral Coordinator Email:		
Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

NURSING

- Provide nursing care per IVX Nursing Procedures, including reaction management and post-procedure observation
NOTE: IVX Adverse Reaction Management Protocol available for review at www.ivxhealth.com/forms (version 09.07.2021)
- TB status and date (Please provide results):

- Hepatitis B status and date (Please provide results):

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- Other: _____
Dose: _____ Route: _____
Frequency: _____

THERAPY ADMINISTRATION

- Certolizumab** (Cimzia) subcutaneous injection
- Induction
 - Dose: 400mg / _____ mg at Week 0, 2, 4, and then with maintenance dosing below
 - Give each 200mg injection separately
- Maintenance
 - Dose: 200mg / 400mg
 - Frequency: every 2 weeks / every 4 weeks / other: _____
- Patient is required to stay for 30-minute observation post infusion/injection
- Patient is NOT required to stay for observation time
- Refills: Zero / for 12 months / _____
(if not indicated order will expire one year from date signed)

SPECIAL INSTRUCTIONS

*Evaluate patients for tuberculosis risk factors and test for latent infection prior to initiating CIMZIA and periodically during therapy. Treatment of latent tuberculosis infection prior to therapy with TNF-blocking agents has been shown to reduce the risk of tuberculosis reactivation during therapy. Prior to initiating CIMZIA, assess if treatment for latent tuberculosis is needed; and consider an induration of 5 mm or greater a positive tuberculin skin test result, even for patients previously vaccinated with Bacille Calmette-Guerin (BCG).

*Test patients for HBV infection before initiating treatment with CIMZIA. For patients who test positive for HBV infection, consultation with a physician with expertise in the treatment of hepatitis B is recommended.

Provider Name (Print)	Provider Signature	Date
-----------------------	--------------------	------

FAX NUMBERS	<input type="checkbox"/> CHICAGO: 312-253-7244	<input type="checkbox"/> INDIANAPOLIS: 844-983-2028	<input type="checkbox"/> PHILADELPHIA: 844-820-9641	<input type="checkbox"/> WEST TN/AR: 888-615-1445
<input type="checkbox"/> BAY AREA: 844-889-0275	<input type="checkbox"/> COLUMBUS: 844-627-2675	<input type="checkbox"/> JACKSONVILLE: 904-212-2338	<input type="checkbox"/> RALEIGH: 919-287-2551	<input type="checkbox"/> MIDDLE TN: 888-615-1445
<input type="checkbox"/> CHARLOTTE: 336-663-0143	<input type="checkbox"/> DAYTONA: 386-259-6096	<input type="checkbox"/> KANSAS CITY: 844-900-1292	<input type="checkbox"/> SARASOTA: 941-870-6550	<input type="checkbox"/> EAST TN: 615-425-7427
<input type="checkbox"/> CINCINNATI: 844-946-0868	<input type="checkbox"/> HARTFORD: 860-955-1532	<input type="checkbox"/> ORLANDO: 844-946-0867	<input type="checkbox"/> SOUTHWEST FL: 813-283-9144	
	<input type="checkbox"/> PALM BEACH: 561-768-9044	<input type="checkbox"/> TAMPA: 844-946-0849		